

Assessment of plant growth and yield of green bean seeds using plasma treatment

*¹Zhuwen Zhou, ²Yanfen Huang and ³Xiaoli He

¹School of physics and electronic science, Guizhou Education University, Guiyang, China

²School of Biological Sciences, Guizhou Education University, Guiyang, China

³Guiyang Baiyun Jinmandi Co., Ltd., Guiyang, Guizhou, China

Corresponding author: stevenzz@foxmail.com

Received on: 03/07/2020

Accepted on: 14/08/2020

Published on: 30/08/2020

ABSTRACT

Aim: The study was carried out to Plasma treatment of seeds at atmospheric pressure as a new technology, through its activation of seed endogenous substances, so that crops improve disease resistance, thereby increasing yield.

Materials and Methods: Using plasma seed processor to treat the seeds of green bean (Jiadowang), the test was carried out with different doses (200 mA~1600 mA).

Results: The results showed that the growth and development of green bean were significantly improved by comparing different plasma doses treated with the untreated CK. and the treatment current (dose) of 1000 mA, 1200mA and 1400 mA of plant traits were significantly changed. The plasma treatment with different doses (current 200 mA~1600mA) can promote the yield of green beans, especially, the yield effect of the appropriate dose (1000 mA) was extremely significant, the plasma current treatments 600 mA, 800mA, 1400mA had a significant effect on the yield of green beans. The trend prediction of single pod yield was basically consistent with the actual test data. The disease resistance of green beans after proper dose (600 mA~1600 mA) plasma treatment was better than that of untreated CK.

Conclusion: In brief, the effect of different doses of plasma treatment on the biological properties of green bean was different. The plant traits of green bean by proper plasma treatment were significantly improved compared with the control (CK), and the disease resistance of treated green bean was also better than that of untreated (CK). The yield of green bean in the eight treated groups could be improved by plasma treatment with different dosage.

Keywords: Green bean, Plasma, Plant traits, Seed, Yield.

How to cite this article: Zhou Z, Huang Y and He X (2020). Assessment of plant growth and yield of green bean seeds using plasma treatment. *J. Agri. Res. Adv.* 02(03): 24-29.

Introduction

At present, physical technology has been widely used in agriculture, plasma treatment of seeds at atmospheric pressure is a new technology, through its activation of seed endogenous substances, so that crops improve stress resistance, thereby increasing yield (Huang and Zhou, 2011). The breeding of traditional crops especially vegetables adopt traditional breeding methods with long cycle and low efficiency. Plasma processing seed technology, is the application of physical methods in agriculture, the cost is lower than the cost of biological agents and chemical agents to treat seeds, and do not pollute the environment (Zhou et al, 2009a and Zhou et al, 2010a).

Copyright: Zhou et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The application of plasma technology in agriculture has a broad prospect. we have used atmospheric pressure dielectric barrier plasma discharge at room temperature for some vegetables (tomato, eggplant, etc.) seed treatment (Zhou et al, 2009b and Zhou et al, 2011a), a decade ago, including the Chinese academy of sciences institute of physics for chili (Chen et al, 2005), lettuce (Wang et al, 2007a), cucumber (Wang et al, 2007b) and other crops to do plasma irradiation technology research, have achieved better results. These techniques mainly use atmospheric pressure dielectric barrier discharge plasma to produce high pressure outside the parallel plate, by adjusting different high pressure to achieve the treatment of different plasma dose of plant seeds, the related vegetable plants have better stress resistance and yield effect. In recent years, high-voltage arc ionization of air nitrogen and oxygen molecules has been used to produce

plasma irradiation combined with alternating inductance, by adjusting the different current treatment of some seeds (Fang et al, 2004 and Geng et al, 2014), but it has been found that the treatment group is not many, the current interval stage is large, the treatment dose is rough, and it is easy to lose some plant biological properties due to the small dose (small current change). In this study, the DL-2 plasma seed processor (Space Machine) was used to treat the seeds before crop sowing. This technology is to draw lessons from the physical effects of space plasma on seeds in space breeding.

In the form of free fall, the seeds have been subjected to short-time light irradiation, inductive action and ozone sterilization to stimulate the potential of seeds, enhance the vigor of seeds and improve the robustness of seeds. After the seeds were treated, the crops had good characters, strong growth and disease resistance, and increased crop yield. therefore, using atmospheric pressure plasma as a new type of seed treatment technology, the combined action of multiple factors will be beneficial to the improvement of varieties, and the technology is carried out at atmospheric pressure under normal temperature without vacuum equipment, easy to operate, low cost, and no damage to plant seed cells. Our research group treated solanaceae seeds (eggplant, tomato) with plasma treatment with different high pressure (4760V~6800V, 4420V~6800V) atmospheric pressure dielectric barrier discharge technology in the original study. On the basis of plasma different currents (200mA~1600mA) were used to treat green bean seeds (Jiadowang), which were tested in nine groups, eight plasma treatment groups and one untreated group (CK) (Ma et al, 2010). The plasma treatment dose was subdivided into 200mA, to observe the changes of plant traits, disease resistance and yield of nine groups. Through mathematical statistics analysis, we can select the best treatment dose of plasma seed treatment to improve the yield of green bean. This research was conducted to provide new technical support for improving yield of green bean and improving efficiency of agricultural production.

Materials and Methods

Materials

The experiments were carried out in the Jinmandi Horticulture Science and Technology Test Field in Penglai Township, Baiyun District, Guiyang

City, Guizhou Province, in March-July 2016, and in the Plasma Laboratory of the Institute of Physics and Electronic Science, Guizhou Education University. The plasma-to-seed treatment device uses a DL-2 plasma seed processor (Space Machine). The test material is green bean (Jiadowang) seeds.

Device

A plasma seed processor (space machine) device DL-2 Dalian Bosi Plasma Co., Ltd was used in this test to treat green bean seeds. The schematic diagram was showed power : <1.2kw; flow rate : >1500kg/h; rated voltage : 220V; frequency : 50Hz; current adjustment range : 0~3.5A; adjustable precision:100mA. When processing, after power on, pull the lower part of the machine air protection switch to the on (ON) position, the voltage indicator pointer swings to the right, showing the supply voltage, up press the plasma switch, the switch is bright, the circuit is on, the plasma emitter in the machine starts, the plasma indication is bright, then the light green light is reflected at the upper feed port of the machine. If the plasma radiant tube does not start, the plasma warning sounds, when the point presses the plasma start button ("point presses" don't hold down, it is pressed, released, pressed again, released again...) until the start of the plasma radiant tube, warning to stop the call. After starting the preheating for 8 min, press the strength switch up to adjust the desired current value by rotating the coarse adjustment and fine-tuning knob. The method of seed treatment is self-flowing, the seed does not stay in the machine, the seed is added from the top feed funnel, and the seed from the outlet must keep the seed flowing smoothly, keep the flowing speed of the seed, and the slow flow rate will affect the effect of seed treatment. The plasma treated seeds were treated once from pouring to receiving, and the number of treatments was 3 times.

Plasma Experiment Method

Plasma seed treatment in a total of 9 groups, 8 treatment groups, and 1 control (CK) group (or untreated group). The seeds were treated with different current by adjusting dosage. The treatment current was 200mA, 400mA, 600mA, 800mA, 1000mA, 1200mA, 1400mA, 1600mA compared with seeds without any treatment (CK). Eight groups were treated three times according to the above dose. It was showed the growth of green beans treated with different doses of

plasma in field (Fig 1) and the growth of green beanun treated group CK (control) in field under the same external environment.

Test Method in Field

Field test design total area was 1 mu (667m²). Line spacing 80cm, plant spacing 50cm, base fertilizer compound fertilizer, the whole growth period topdressing 2 times, leaf surface fertilizer 3 times and with headache powder, amoxicillin pest control. Only a small amount of leaf surface rust spots, through leaf mesh headache powder, can be controlled. Plant traits, yield traits, disease resistance and growth period were recorded.

Results and Discussion

Effects of Plasma Treatment with Different Dose (Current) on Plant Traits of Green Beans

Effects on the Stem coarse of Green bean

Effects on the stem coarse of green bean from the plant traits showed that the plasma treatment of 8 groups increased (0.2~0.5 cm) compared with untreated group CK (Table 1). Treatment groups No. 5,6,7,8 increased significantly, the other treatment groups were 0.3 cm. higher than the untreated CK. Zhou et al, (2011b) also corroborated with the findings of this study.

Effects on the Number of Branches of Green beans

After plasma treatment, the effect of branch number of 8 groups was that there was no change in treatment groups No.2,3 and 8, the number of branches of treatment groups No.4,5,6 and 7 increased by 1 (contrast CK), but the number of branches of treatment group No.1 decreased by 1. These findings were in agreement with results of Chen et al (2005).

Effect on the Node position of the First inflorescence of Green bean

Node position of the first inflorescence of the plasma treatment groups were affected, For the treatment groups No. 1, 3, 5, their first inflorescence was decreased compared with CK group, no change in other treatment groups according to Wang et al (2007b)

Effects on the Fruit number of per Inflorescence of Green beans

Effect of the fruits of inflorescence per inflorescence in the eight groups treated by plasma, except for treatment group No.1, the fruit number of per inflorescence of the other treatment groups increased by 2 than the control CK. The results of the study were in line with Zhou et al, (2010c)

Effect on the Length of Single pod in Green bean

Except for no change in treatment No.1, the single pod length of the seven groups treated by plasma increased to varying degrees compared with the control CK (1cm~4cm), the longer increasing was No.4~No.8 groups.

To sum up, different dose plasma treatment of green bean seeds, resulting in different degrees of impact on green bean stem coarse, number of branches, node position of the first inflorescence, of fruits per inflorescence, single pod length plant traits, and each botanical character index of number different dose plasma treatment is also different, most indicators change significantly, a few are not obvious or no change. All in all, the growth and development of green beans were obviously promoted by proper plasma treatment at different doses (compared with CK), especially the treatment group with significant changes in the influence of green bean plant traits was the No.5, No.6 and No.7 groups in agreement Zhou et al, (2010b)

Effect of Plasma Treatment with Different Dose (Current) on Disease Resistance of Green Bean

Disease resistance: compared with untreated (CK), no viral rust spots was found in six groups (No. 3,4,5,6,7,8) treated by plasma except treatment groups No. 1,2 and control group CK (Table 1). The reason may be due to insufficient plasma treatment dose in treatment groups No. 1 and 2. All nine groups (8 treatment groups and 1 CK group) had no leaf mold and powdery mildew. The fertility period of the other 6 treatment groups (groups No. 3,4,5,6,7,8) was 5~10 days longer than that of the CK group, except for the first, second and CK groups, which had about 100 days. Overall, proper dose (600mA~1600mA) plasma treatment showed better resistance to disease than untreated CK which was in line with Huang and Zhou (2011).

Effects of different doses of plasma treatment on yield traits of green bean

The yield traits (Table 1): Compared with untreated (CK), the yield of single pod in all 8 groups treated by plasma was higher than that in control (CK), and the yield of single pod in 8 groups increased 1kg~3kg. The first two highest yield were: treatment group No. 5 single pod yield 9 kg~13kg (plasma dose 1000mA), treatment group No. 7 pod yield 8 kg-13kg (plasma dose 1400mA), corresponding to single pod yield 28.57%~30%, 14.28%~30% higher than CK, respectively. Lowest single pod yield was also increased by 14.28% (plasma dose 200mA).

From the start-up period, the start-up period of all 8 groups treated by plasma was 2~6 days more than that of untreated CK. To sum up, plasma treatment with different doses (plasma current 200mA~1600mA) can promote the growth, development and yield of green bean, in which the treatment yield effect of appropriate dose (1000mA) was extremely significant, and the plasma current treatment 600mA, 800mA, 1400mA had a significant effect on green bean yield according to Zhou et al (2011b).



Fig 1. (Color online) The growth of green beans treated with different doses of plasma in field.




Fig 2. (Color online) Mathematical statistics of yield trend of green bean by plasma treatment with different doses.

Table1: Record of effects of plasma treatment on Green Bean

Groups No.	Dose (mA)	Plant traits				Yield traits			Pest/plant		Fertility traits	
		Stem coarse (cm)	Number of branches (each)	Node position of first inflorescence (each)	Fruit number per Inflorescence (each)	Single pod length (cm)	Yield (kg)	initial collection period (day)	viral rust spots	leaf mold	powdery mildew	growth stage (day)
1	200	0.8~1.5	3~4	4~5	2~6	12~16	8~10	38	√			90~100
2	400	0.8~1.5	3~5	4~6	4~6	13~17	8~11	40	√			100
3	600	0.8~1.5	3~5	4~5	4~6	14~18	9~12	40				100~105
4	800	0.8~1.5	3~6	4~6	4~6	14~19	9~12	42				100~105
5	1000	0.8~2.0	3~6	4~5	4~6	13~20	9~13	40				100~110
6	1200	0.8~2.0	3~6	4~6	4~6	13~19	8~12	40				100~110
7	1400	0.8~2.0	3~6	4~6	4~6	13~19	8~13	40				100~106
8	1600	0.8~2.0	3~5	4~6	4~6	13~19	8~12	40				100~106
9	0	0.5~1.5	3~5	4~6	4	12~16	7~10	36	√			100

Mathematical and statistical analysis of yield (Series 1) trend of green bean single pod

The results showed that the yield of single pod was predicted from the regression trend line (Fig 2). The relationship between single pod yield and plasma current treatment (regression equation) was in accordance with the two-period average moving distribution. The first peak appeared at (1000mA) and the second at (1400mA). The results showed that the two treatment groups (groups No. 5 and 7) with the selected plasma dose of 1000mA and 1400mA were the most productive for single pod. The trend prediction of single pod yield is basically consistent with the actual test data. Zhou et al, (2011a) and Geng et al (2014) also advocated the findings of the study.

Mathematical and statistical analysis of yield (Series 2) trend of green bean single pod

The mathematical statistical analysis of yield of single pod (series 2) was performed after the treatment of unequal ion doses (Fig 2): prediction of green bean from regression trend line: the relationship between yield of single pod and plasma current treatment (regression equation) was basically in accordance with the Gaussian distribution law, which is consistent with our previous research prediction (Zhou et al, 2011a). The yield of green bean was greatly improved after plasma treatment of seeds with an appropriate plasma dose range of 600mA~1000 mA (groups No. 3, 4, 5). The peak of Gaussian distribution prediction appears around

800mA~1000 mA, and the yield trend prediction was basically consistent with the actual test data. These findings were in agreement with the results of Fang et al (2004).

Conclusion

It is undoubtedly of great economic and strategic significance to use high voltage arc ionization air nitrogen and oxygen molecules to produce plasma irradiation and alternating inductance to treat plant seeds by adjusting different current (dose), to change their growth law, to improve product quality and to increase local crop yield. This kind of applied research has a good prospect for solving some key technical problems in national economy and social development. It can provide a possible and effective way for more crops to increase production, and also can build a technical platform for the improvement of traditional agricultural breeding technology means and the application of high and new technology in agricultural breeding. In particular, the application of plasma technology to agricultural science, biological science and other interdisciplinary development has far-reaching scientific significance. The effects of plasma treatment on the biological properties and yield of green bean (Jiadouwang) were studied by using plasma seed processor and different dose (200mA~1600mA). The results showed that the plasma irradiation combined with alternating inductance was used to induce the physiological changes of the seeds.

The effects of physiological changes on the yield traits of the plants were investigated and analyzed, such as stem coarse, branch number, first inflorescence node position, fruit number of inflorescence, single pod length, and seed disease resistance. The results showed that different current (dose) plasma treatment had different effects on the plant traits of green bean. The growth and development of green beans were obviously promoted by proper plasma treatment at different doses, especially the treatment current (dose) with a significant change in the effect on the plant traits of green bean was 1000mA, 1200mA and 1400mA (treatment groups were No. 5, 6, 7), respectively. The plasma treatment with different doses (plasma current 200mA~1600mA) can promote the growth, development and yield of green bean. The treatment yield effect of the appropriate dose (1000mA) is extremely significant. The plasma

current treatment 600mA, 800mA, 1400mA has a significant effect on green bean yield. The trend prediction of single pod yield is basically consistent with the actual test data. The disease resistance of green beans after proper dose (600mA~1600mA) plasma treatment was better than that of untreated CK.

All in all, the effect of different doses of plasma treatment on the biological properties of green bean was different. The plant traits of green bean by proper plasma treatment were significantly improved compared with the control (CK), and the disease resistance of treated green bean was also better than that of untreated (CK). The yield of green bean in the eight treated groups could be improved by plasma treatment with different dosage.

Acknowledgments

This project was supported by the Natural Science Research Project of the Key Laboratory of Guizhou Province Education Bureau Grant No. KY[2014]217, of the Scientific Research Innovation team in plasma and functional thin film materials of Guizhou Province Education Bureau Grant No. QJTD[2014]38, of Guizhou Province Key Support Discipline "Theoretical Physics" Grant No. ZDXK[2015] 38, of Guizhou Province First-class Major-Physics Grant No. JGTH[2019] 46, of Key Subjects at School Level-Plasma Physics, of Institute of Plasma Physics Applications.

References

- Chen GL, Fan SH, Li CL, Gu WC, Feng WR, Zhang GL, Wang JL, Latif K, Zhang SG, Wang ZQ, Han EL, Fu YB, Yang SZ (2005). A novel atmospheric pressure plasma fluidized bed and its application in mutation of plant seed. *China Physics Letter*, 22 (8): 1980-1983.
- Fang XQ, Bian SF and Xu KZ (2004). Effects of Plasma Treatment on Biological Properties and Yield of Maize Seeds. *Journal of Maize sciences*, 12 (4): 60-61.
- Geng SJ, Geng LZ and Chen ZG (2014). Effect of Plasma Seed Processing Machine on Maize. *Modern agriculture*, (2): 28-29.
- Huang YF and Zhou ZW (2011). Effect of atmospheric pressure plasma technology on eggplant. *Seeds*, 29 (1): 73-75.
- Ma XL, Qu QH and Liu F (2010). Effect of Plasma Seed Processing Machine on Maize. *Modern agriculture*, (5): 15-16.

- Wang M, Chen QY and Yang SZ (2007a). Effects of atmospheric pressure plasma treatment on seed germination and growth of lettuce. *North China Journal of Agriculture*, 22 (6): 108-113.
- Wang M, Yang SZ and Chen QY (2007b). Effects of atmospheric pressure plasma treatment on seed germination and seedling growth of cucumber. *Journal of Agricultural Engineering*, 23 (2): 195-200.
- Zhou ZW, Huang YF, Chen W and Yang SZ (2011a). A New Atmospheric Pressure Plasma Equipment and Its Application in Eggplant Seeds. *Guizhou Agricultural Science*, 39 (3): 61-65.
- Zhou ZW, Huang YF, Deng MS and Yang SZ (2009a). Effects of Atmospheric Pressure Plasma Treatment on Yield and Quality of Eggplant of No.2-Qianqie Seed. *Modern Agricultural Technology*, 19: 94-96.
- Zhou ZW, Huang YF, Yang SZ and Chen W (2011b). Introduction of a new Atmospheric Pressure Plasma Device and Application on Tomato Seeds. *Agricultural Sciences*, 2 (1): 23-27.
- Zhou ZW, Huang YF, Yang SZ and Deng MS (2009b). Effects of Plasma Treatment on Plant Growth Development and Yield Quality of No .10-Hongza Seed. *Guizhou Agricultural Science*, 37 (12): 58-61.
- Zhou ZW, Huang YF, Yang SZ and Deng MS (2010a). Effect of atmospheric pressure plasma treatment on growth, development, yield and quality of tomato. *Anhui Agricultural Science*, 38 (2): 1085-1088.
- Zhou ZW, Huang YF, Yang SZ and Deng MS (2010b). Effect of Atmospheric Pressure Plasma on Plant Growth and Yield of Eggplant. *Chinese Vegetables*, (4):62-66.
- Zhou ZW, Yang SZ, Huang YF and Chen W (2010c). Effects of Atmospheric Pressure Plasma on Growth and Development of Tomato. *International Conference on Agricultural Sciences and Engineering*, Macau, China, 10-11 September. 55-57.
